

Research Article



## INFLUENCE OF BIOACTIVE SPICE EXTRACTS ON THE FUNCTIONAL AND MICROBIAL INTEGRITY OF A TRADITIONAL FERMENTED RICE FOOD

Masauda Muhammad Umar<sup>1\*</sup>, Innocent Ojeba Musa<sup>2</sup>, Miracle Uwa Livinus<sup>3</sup>,  
Mustapha Abdulsalam<sup>4</sup>, Tasnim Tasnim<sup>5</sup>, Firdaus Fahdi<sup>6</sup>, Amalia Riski Wahyulita<sup>7</sup>

<sup>1,2,4</sup>Department of Microbiology, Skyline University Nigeria

<sup>3</sup>Department of Biochemistry, Skyline University Nigeria

<sup>5,6,7</sup>Deli Husada Deli Tua Health Institute, Indonesia

**Corresponding Author :**

Innocent Ojeba Musa

e-mail: [innocentmusa0011@gmail.com](mailto:innocentmusa0011@gmail.com)<sup>2\*</sup>

### ABSTRACT

**Background:**

This study investigates the impact of incorporating natural spices ginger (*Zingiber officinale*), garlic (*Allium sativum*), and cloves (*Syzygium aromaticum*) at concentrations of 1%, 3%, and 5% on the microbial stability, physicochemical properties, nutritional composition, sensory attributes, and safety of Sinasir (Rice pancake), a traditional Nigerian fermented rice cake.

**Methods:** Over an 8-week storage period, samples were analyzed for microbial load, pH, temperature, titratable acidity, proximate composition (carbohydrates, moisture, fat, ash, protein, and fiber), sensory evaluation, hazard analysis, and antimicrobial sensitivity.

**Results:** this finding indicated that spice-treated samples, particularly those with 5% clove and garlic, exhibited significantly lower bacterial and coliform counts compared to the control, which reached TNTC by week 4. The 5% clove sample maintained bacterial counts below  $3.6 \times 10^5$  CFU/g at week 8. pH levels in spiced samples remained acidic (4.2–4.5), while the control increased to 5.8, correlating with spoilage. Moisture content decreased with higher spice concentrations, enhancing shelf stability. The 5% garlic sample showed the highest protein content (9.85%), and the 5% clove sample had the highest ash content (2.41%). Sensory evaluation favored 3% and 5% spice-treated samples. The bacterial isolates include *Bacillus cereus*, *Pseudomonas aeruginosa*, *Enterobacter aerogenes*, and *Lactobacillus fermentum*. Antimicrobial assays revealed significant inhibition zones against foodborne pathogens, particularly for clove and garlic extracts.

**Conclusion:** These findings support the use of local spices in improving the safety, quality, and acceptability of traditional fermented foods.

**Keywords:** Sinasir, Rice pancake, ginger, garlic, cloves, fermented food, microbial load, proximate, analysis, antimicrobial, activity, sensory, quality

## INTRODUCTION

Food is more than sustenance; it is memory, culture, and identity. Across the world, fermented foods occupy a unique space at this intersection. They are celebrated not only for their flavors and textures but also for their ability to transform simple ingredients into nutritionally enhanced, microbiologically diverse, and shelf-stable products. From kimchi in Korea to sauerkraut in Germany, injera in Ethiopia, and Sinasir in Northern Nigeria, fermentation embodies both ancient ingenuity and modern relevance [1]. In resource-limited settings, fermented foods remain indispensable, offering affordable nutrition to millions who face food insecurity [2].

Despite their importance, traditional fermented foods are vulnerable. Their production often occurs under informal conditions such as family kitchens, community gatherings, or small-scale roadside vendors where temperature control, hygienic monitoring, and microbial surveillance are difficult to enforce. Consequently, the safety of such foods is frequently compromised by contamination with pathogenic microorganisms. Outbreaks of foodborne illness continue to underscore this vulnerability, raising critical questions about how to preserve the cultural integrity of these foods while ensuring their safety and marketability in a globalized world [3].

One promising solution lies within the same cultural frameworks that gave rise to these foods in the first place: the use of spices. For centuries, communities have incorporated spices not only for flavor but also for preservation. Garlic, ginger, and cloves are ubiquitous in African and Asian cuisines, valued in both kitchens and traditional medicine. Modern science affirms

what ancient cultures intuited. These spices harbor bioactive compounds such as allicin in garlic, gingerol in ginger, and eugenol in cloves that confer antimicrobial and antioxidant effects [4]. By integrating these naturally occurring compounds into fermentation processes, it may be possible to mitigate microbial risks while enhancing nutritional and sensory quality.

The case of Sinasir, a rice-based pancake widely consumed in Northern Nigeria, illustrates this challenge and opportunity. Beloved for its soft texture and mild sour flavor, Sinasir is more than a meal; it is a staple of community life, served during festivals, weddings, and family gatherings. Yet its short shelf life and susceptibility to microbial spoilage limit its distribution beyond local markets [5]. Without intervention, its role risks being diminished in an era where food safety standards increasingly dictate global trade and acceptance. By applying locally accessible spice extracts, it may be possible to preserve not only the food but also the cultural continuity it represents.

This endeavor has broader implications. The growing global movement toward “clean label” foods, which refers to products free of synthetic preservatives, demands natural and sustainable solutions [6]. Fermented foods enhanced with bioactive spices align with this demand by bridging traditional knowledge with contemporary food science. Moreover, with climate change threatening food supply chains and antimicrobial resistance rising, the search for natural preservation strategies is more urgent than ever [7].

In this study, we investigate how spice extracts of garlic, ginger, and clove affect the microbial stability, nutritional composition, and sensory attributes of Sinasir. Beyond the

laboratory, the aim is to demonstrate how traditional ethnobotanical practices can be harnessed to address modern food security challenges. By situating local foods within a global scientific framework, this study highlights how indigenous knowledge remains a reservoir of solutions for future food sustainability.

## MATERIAL AND METHODS

### Sample Collection and Preparation

The core ingredients for Sinasir (rice pancake), namely rice, sugar, yeast, and salt, were purchased from Kurmi Market in Kano, Nigeria. Preparation followed traditional household practices with controlled adjustments to ensure reproducibility. A total of 600 g of rice was used. Out of this, 550 g was soaked in water for 12 hours to initiate partial fermentation, while 50 g was cooked to provide gelatinized starch that improves texture. The soaked and cooked rice were then combined and wet-milled into a batter. Yeast, sugar, and salt were added in standardized quantities, after which the mixture was allowed to ferment for 5 hours. The fermented batter was stirred to incorporate air, reflecting artisanal preparation techniques, and pan-fried to produce Sinasir.

### Preparation of Spice Extracts

Ginger (*Zingiber officinale*), garlic (*Allium sativum*), and cloves (*Syzygium aromaticum*) were chosen because of their widespread culinary and medicinal use in Nigeria. After thorough washing, the spices were sliced, shade-dried, and ground into fine powders. Fifty grams of each spice were soaked in 120 ml of distilled water at room temperature for 72 hours to extract bioactive compounds, following the method described by Sah et al. 2020 [8]. The extracts were

filtered through Whatman filter paper to remove solid particles and refrigerated at 4°C until use. To evaluate concentration-dependent effects, spice extracts were incorporated into the Sinasir batter at 1 percent, 3 percent, and 5 percent. Prepared pancakes were labeled, refrigerated, and observed over an 8-week storage period.

### Microbial Enumeration and Isolation

Microbial stability was evaluated by determining total bacterial counts immediately after production and at weekly intervals during storage. Twenty-five grams of each sample were aseptically homogenized in 225ml of buffered peptone water [9]. Serial dilutions were prepared and plated using the pour-plate technique. One milliliter of each dilution was transferred into sterile Petri dishes and mixed with molten nutrient agar maintained at 45°C. The plates were gently swirled to ensure uniform distribution, solidified, and incubated at 37°C for 24 hours. Plates with 30 to 300 colonies were enumerated, and the results expressed as colony-forming units per gram (CFU/g) [10]. Fungal counts were assessed using the same protocol, except that Potato Dextrose Agar (PDA) was used and plates were incubated at room temperature for 3 to 5 days.

### Microbiological Characterization

Standard plate count methods were used to evaluate viable bacteria, coliforms, and fungi at weeks 0, 2, 4, 6, and 8 [11]. Isolates were further characterized using biochemical tests including catalase, citrate, methyl red–Voges Proskauer (MR-VP), indole, and triple sugar iron (TSI). These tests provide insights into the metabolic activities of microorganisms [12].

## Physicochemical Analyses

### *pH Measurement*

The acidity of Sinasir was monitored as a marker of fermentation. Three grams of sample were homogenized in 10 ml of distilled water. The slurry was transferred into a beaker, and a calibrated pH meter (Model 501) with a temperature probe was immersed until a stable reading was obtained [13].

### *Titrateable Acidity*

Titrateable acidity was determined as described by Akpakpan et al [14]. A 3 g sample was homogenized in 10 ml of distilled water, after which five drops of phenolphthalein indicator were added. The mixture was titrated against 0.1 N NaOH until the first permanent pink coloration was observed. Results were expressed as percentage lactic acid equivalents, and ambient temperature (25–28°C) was recorded during analysis.

### *Proximate Composition*

Nutritional quality was determined by proximate analysis of moisture, ash, protein, fat, crude fiber, and carbohydrate contents using standard methods [11].

### *Sensory Evaluation*

To capture consumer perception, sensory evaluation was conducted by a trained panel of 20 members. Samples coded to maintain blinding were evaluated for color, taste, aroma, texture, and overall acceptability using a 9-point hedonic scale [15].

### *Hazard Analysis*

Hazard Analysis and Critical Control Points (HACCP) was applied to identify stages in Sinasir preparation that posed food

safety risks. Special attention was given to raw material sourcing, fermentation, and cooking, which were considered critical points for microbial control.

### *Antimicrobial Sensitivity Testing*

The antibacterial activity of spice extracts was tested against selected foodborne pathogens including *Escherichia coli*, *Lactobacillus fermentum*, *Enterobacter aerogenes*, *Bacillus cereus*, and *Pseudomonas* spp. Agar well diffusion was used as the assay method. After 24 hours of incubation, inhibition zones were measured in millimeters to quantify antimicrobial effectiveness [16].

## RESULTS

### *Microbial Stability of Sinasir During Storage*

Spice-treated samples exhibited significantly lower bacterial, coliform, and fungal loads compared to the untreated control (Fig. 1–3). While microbial proliferation increased over time in all samples, higher spice concentrations (particularly 5% garlic and 5% clove) delayed spoilage, with counts remaining within acceptable food safety limits up to week 6. In contrast, control samples exceeded countable microbial loads as early as week 3, underscoring the vulnerability of Sinasir without preservation.

These findings confirm the antimicrobial potential of traditional spices. Garlic's allicin and clove's eugenol showed the strongest inhibition, consistent with earlier reports of their broad-spectrum activity [17]. Ginger, while moderately effective, showed reduced control over extended storage, suggesting its compounds degrade more rapidly.




Fig.1. Effect of different spice treatment on bacterial count of Sinasir (Rice pancake) at different concentration.




Fig.2. Effect of different spice treatment on coliform count of Sinasir (Rice pancake) at different concentration.




Fig.3. Effect of different spice treatment on Fungal count of Sinasir (Rice pancake) at different concentration

**Physicochemical Properties**

Changes in pH, titratable acidity, and temperature reflected microbial activity and preservation effects. Spice-treated samples maintained greater stability compared to the

control. Notably, clove-treated samples exhibited the least pH fluctuation across storage, highlighting its capacity to moderate fermentation dynamics (Fig. 4–6).




Fig.4. Effects of various treatments on the Titratable acidity of Sinasir (Rice pancake) at different concentration




Fig. 5. Effects of various treatments on the Temperature of Sinasir (Rice pancake) samples at different concentration.




Fig.6. Effects of various treatments on the pH of Sinasir (Rice pancake) at different concentration

### Nutrient Composition

Proximate analysis revealed gradual declines in protein, fiber, fat, and carbohydrates during storage (Fig. 7–12). However, garlic and clove at higher concentrations mitigated these losses, likely

due to antioxidant activity that slowed degradation. Moisture levels also declined steadily, with spice-treated samples showing more controlled reduction, preserving textural integrity.




Fig.7. Effects of various treatments on the Ash content of Sinasir (Rice pancake) samples at different concentration.




Fig.8. Effects of various treatments on the Moisture content of Sinasir (Rice pancake) at different concentration.




Fig.9. Effects of various treatments on the Crude protein of Sinasir (Rice pancake) at different concentration.




Fig.10. Effects of various treatments on the Crude fiber of Sinasir (Rice pancake) at different concentration




Fig.11. Effects of various treatments on the Fat content of Sinasir (Rice pancake) at different concentration.




Fig.12. Effects of various treatments on the Carbohydrates of Sinasir (Rice pancake) at different concentration.

### Sensory Evaluation

Sensory scores (taste, aroma, texture, color, and overall acceptability) favored spice-treated samples, with garlic- and clove-enriched Sinasir ranking highest in taste and

aroma (Fig. 13). Importantly, spices did not diminish visual appeal or textural qualities, suggesting they can enhance both shelf-life and consumer acceptance.




Fig.13. Mean sensory scores of Sinasir (Rice pancake) at different concentration

### Hazard Analysis

HACCP evaluation identified soaking and cooling as critical points of microbial risk. However, frying effectively eliminated detectable microbial loads (Fig. 14). The reappearance of microbes after cooling

highlights the importance of hygienic handling post-cooking. Spices provided additional microbial suppression, but strict handling practices remain indispensable.




Fig.14. Microbial load at different stages of Sinasir (Rice pancake) product

### Antimicrobial Sensitivity

Aqueous spice extracts displayed concentration-dependent antibacterial

activity (Tables 2–4). Garlic was the most potent, producing inhibition zones up to 17.3 mm against *Enterobacter aerogenes*. Ginger

demonstrated moderate activity, while clove showed limited efficacy in aqueous form, reflecting eugenol’s low solubility in water. These results reinforce the importance of extraction method optimization for maximizing antimicrobial potential.

**Table 1: Results of Antibacterial Effect of Aqueous Extract of *Allium sativum* on the test organisms**

Conc. Mg/ml	<i>Pseudomonas Aeruginosa</i>	<i>Bacillus Cereus</i>	<i>Lactobacillus fermentum</i>	<i>Enterobacter Aerogenes</i>
0.2	2.9mm	4.2mm	4.9mm	3.5mm
0.4	5mm	5.1mm	7.5mm	6.8mm
0.6	13mm	10.2mm	6.4mm	4.3mm
0.8	9mm	7.1mm	8.2mm	11.9mm
1.0	16mm	12.8mm	10.8mm	17.3mm

**Table 2: Results of Antibacterial Effect of Aqueous Extract of *Syzgium aromaticum* on the test organisms**

Conc. Mg/ml	<i>Pseudomonas aeruginosa</i>	<i>Bacillus Cereus</i>	<i>Lactobacillus fermentum</i>	<i>Enterobacter Aerogenes</i>
0.2	NA	NA	NA	NA
0.4	NA	NA	NA	NA
0.6	NA	NA	NA	NA
0.8	0.6mm	NA	NA	NA
1.0	1.5mm	1.1mm	NA	2.0mm

NA-Not detected

**Table 3: Results of Antibacterial Effect of Aqueous Extract of *Zingiber officinale* on the test organisms**

Conc. Mg/ml	<i>Pseudomonas Aeruginosa</i>	<i>Bacillus Cereus</i>	<i>Lactobacillus fermentum</i>	<i>Enterobacter Aerogenes</i>
0.2	NA	NA	NA	NA
0.4	NA	1.7mm	NA	1.5mm
0.6	3.8mm	3.9mm	3.3mm	2.9mm
0.8	4.5mm	5.00mm	5.1mm	4.0mm
1.0	6.2mm	8.1mm	7.9mm	6.5mm

NA-Not detected

## DISCUSSION

The results of this study demonstrate that incorporating spice extracts, particularly garlic and cloves, significantly reduced bacterial, coliform, and fungal loads in Sinasir during storage, while also preserving nutritional and sensory qualities. These

findings are in strong agreement with Adeyanju et al. (2022) [18] and Hassan et al. (2020) [19], who emphasized the antimicrobial efficacy of traditional spices in extending the shelf life of African fermented foods. Similarly, the dose-dependent effects of ginger, garlic, and cloves observed here

align with reports by Gholami-Shabani et al. (2017) [20] and Gholipour Kanani et al. (2014) [21], who showed that higher spice concentrations delay microbial proliferation more effectively. The persistence of *Bacillus cereus* and *Pseudomonas aeruginosa* despite treatment reflects earlier observations by Omotayo et al. (2020) [22] and Yehia et al. (2022) [23] that spore-formers and adaptive spoilage organisms often resist natural antimicrobials, suggesting that spices alone may not provide complete inhibition. Nutritional stability, particularly in protein and ash content, corroborates the findings of Bou et al. (2009) [24] and Ozma et al. (2003) [25], who noted that bioactive compounds in garlic and cloves preserve structural integrity and slow nutrient degradation. However, the relatively weaker and inconsistent antimicrobial effect of ginger diverges from studies such as Žitek et al. (2022) [26], where ginger extracts showed stronger inhibition under controlled conditions, indicating that food matrix complexity and storage conditions may influence efficacy.

Collectively, this study both confirms and nuances earlier research by showing that spices enhance the microbial safety and consumer acceptability of traditional foods, but their effectiveness may depend on concentration, food composition, and storage duration [27].

## CONCLUSION

This study examined the role of ginger, garlic, and clove extracts in shaping the microbial, nutritional, physicochemical, and sensory characteristics of Sinasir, a traditional Nigerian rice-based pancake that holds deep cultural significance. By integrating these natural spices into the fermentation process, we demonstrated that food safety, nutritional stability, and

consumer acceptability can be significantly improved without compromising authenticity.

The findings reveal that garlic and clove, in particular, exhibit strong antimicrobial effects that slowed spoilage and maintained microbial loads within safe limits during extended storage. These spices also helped preserve critical nutrients such as protein and minerals while moderating physicochemical changes in pH and titratable acidity, which are often linked to product deterioration. From a sensory perspective, panelists consistently rated spice-treated samples as acceptable and in some cases even preferable, underscoring the potential of these natural additives to enhance both shelf-life and consumer enjoyment. Importantly, hazard analysis confirmed that the integration of spices into production can mitigate risks at critical control points, offering an added layer of protection in contexts where modern food preservation systems are often inaccessible.

Beyond the laboratory, the implications of these findings are substantial. In Nigeria and across sub-Saharan Africa, where food insecurity and foodborne illnesses remain pressing public health challenges, culturally embedded strategies like the use of indigenous spices represent practical and sustainable solutions. They provide households and small-scale vendors with affordable tools to safeguard traditional foods while aligning with global calls for clean-label, chemical-free preservation. On a broader scale, this work highlights the untapped potential of ethnobotanical knowledge in addressing modern scientific and societal concerns, from antimicrobial resistance to sustainable food systems.

In conclusion, the incorporation of garlic, clove, and ginger into Sinasir

production is more than a technical adjustment; it represents a bridge between tradition and innovation. By validating long-standing culinary practices through rigorous scientific analysis, this study contributes to the preservation of cultural heritage while offering a pathway toward safer, more nutritious, and more resilient food systems. Future research should build on these findings by exploring synergistic combinations of spices, optimizing extraction methods, and scaling applications to other fermented foods within and beyond Africa. In doing so, local traditions may inspire global innovations in food safety and sustainability.

## REFERENCES

1. Marco, M. L., Sanders, M. E., Gänzle, M., Arrieta, M. C., Cotter, P. D., De Vuyst, L., ... & Hutkins, R. (2021). The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on fermented foods. *Nature Reviews Gastroenterology & Hepatology*, 18(3), 196-208.
2. FAO. (2021). *Food Safety in Africa: Past, Current and Future*. Food and Agriculture Organization of the United Nations.
3. Bell, V., Guina, J., & Fernandes, T. H. (2023). African fermented foods and beverages. Potential impact on health. *Microbial fermentations in nature and as designed processes*, 293-322.
4. Avci, G. A., Avci, E., Cilak, G. O., & Cevher, S. C. (2020). Antimicrobial and Antioxidant Activities of Zingiber officinale Ginger and Alpinia officinarum Galangal. *Hittite Journal of Science and Engineering*, 7(1), 45-49.
5. Olaniran, A. F., & Abiose, S. H. (2022). Fermented rice products in Nigeria: Microbiology, safety, and prospects. *African Journal of Microbiology Research*, 16(2), 67–75.
6. Chauhan, A., & Singh, S. K. (2022). Antimicrobial properties of spices: A review. *Journal of Food Science and Technology*, 59(3), 1234–1245.
7. WHO. (2020). *Global Strategy for Food Safety 2020–2030*. World Health Organization.
8. Sah, S. N., Khanal, H., & Acharya, D. R. (2020). Antibacterial activity of common spices extracts on bacterial isolates found in Kachhila, a Newari cuisine. *Tribhuvan University Journal of Microbiology*, 7, 8-18.
9. Vasavada, P. C., Lee, A., & Betts, R. (2020). Conventional and novel rapid methods for detection and enumeration of microorganisms. In *Food safety engineering* (pp. 85-128). Cham: Springer International Publishing.
10. Yousef, A. E., & Carlstrom, C. (2003). *Food microbiology: A laboratory manual*. John Wiley & Sons.
11. AOAC. (2005). *Official Methods of Analysis* (18th ed.). Association of Official Analytical Chemists.
12. Cheesbrough, M. (2005). *District laboratory practice in tropical countries, part 2*. Cambridge university press.
13. Horwitz, W. J. G. B. C. (1975). *Association of official analytical chemists (AOAC) methods*. George Banta Company, Menasha, WI.
14. Akpakpan, A. E., Uwanta, E. J., Akpan, P. E., & Owen, A. E. (2020). Phytochemical screening and analytical assessment of acid-base indicator properties of red and white kola nuts

- extracts. *Asian Journal of Advances in Research*, 3(1), 1-7.
15. Bamidele, F. S., Adeyanju, M. M., & Okonkwo, C. (2023). Sensory perspectives into indigenous fermented foods in the tropics. *African Journal of Food Science and Technology*, 14(2), 45–56.
  16. Chauhan, K., & Rao, A. (2024). Clean-label alternatives for food preservation: An emerging trend. *Heliyon*, 10(16).
  17. Sarfraz, M., Nasim, M. J., Jacob, C., & Gruhlke, M. C. (2020). Efficacy of allicin against plant pathogenic fungi and unveiling the underlying mode of action employing yeast based chemogenetic profiling approach. *Applied Sciences*, 10(7), 2563.
  18. Adeyanju, J. A., Ogunlakin, G. O., Abioye, A. O., Alabi, O. D., Oloyede, A. A., & Akinola, F. B. (2022). Assessment of biscuits produced from the blends of wheat, acha and germinated African yam bean flour. *Journal of Advances in Food Science & Technology*, 9(2), 26-37.
  19. Hassan, M. Z., Malgwi, J. I., Usman, A., Barde, Z. M., Abdullahi, A. I., & Usman, U. (2023). Effect of Prolonged Storage in Ice on Nutrients Composition and Sensory Quality of Fresh Pond Raised Catfish (*Clarias gariepinus* Burchell, 1822). *FUDMA JOURNAL OF SCIENCES*, 7(3), 288-292.
  20. Gholami-Shabani, M., Gholami-Shabani, Z., Shams-Ghahfarokhi, M., Jamzivar, F., & Razzaghi-Abyaneh, M. (2017). Green nanotechnology: biomimetic synthesis of metal nanoparticles using plants and their application in agriculture and forestry. In *Nanotechnology: An Agricultural Paradigm* (pp. 133-175). Singapore: Springer Singapore.
  21. Gholipour Kanani, H., Nobahar, Z., Kakoolaki, S., & Jafarian, H. (2014). Effect of ginger-and garlic-supplemented diet on growth performance, some hematological parameters and immune responses in juvenile *Huso huso*. *Fish physiology and biochemistry*, 40(2), 481-490.
  22. Omotayo, A. O., & Aremu, A. O. (2020). Underutilized African indigenous fruit trees and food–nutrition security: Opportunities, challenges, and prospects. *Food and Energy Security*, 9(3), e220.
  23. Yehia, H. M., Al-Masoud, A. H., Elkhadragey, M. F., Sonbol, H., & Al-Dagal, M. M. (2022). Analysis of spore-forming bacterial contaminants in herbs and spices and evaluation of their heat resistance. *Food Science and Technology*, 42, e19422.
  24. Bou, R., Codony, R., Tres, A., Decker, E. A., & Guardiola, F. (2009). Dietary strategies to improve nutritional value, oxidative stability, and sensory properties of poultry products. *Critical reviews in food science and nutrition*, 49(9), 800-822.
  25. Ozma, M. A., Abbasi, A., Ahangarzadeh Rezaee, M., Hosseini, H., Hosseinzadeh, N., Sabahi, S., ... & Kafil, H. S. (2023). A critical review on the nutritional and medicinal profiles of garlic's (*Allium sativum* L.) bioactive compounds. *Food Reviews International*, 39(9), 6324-6361.
  26. Žitek, T., Kučuk, N., Postružnik, V., Leitgeb, M., Knez, Ž., Primožič, M., & Marevci, M. K. (2022). Synergistic effect of supercritical and ultrasound-

assisted ginger (*Zingiber officinale* Roscoe) extracts. *Plants*, 11(21), 2872.

27. Innocent Ojeba Musa, Job Oloruntoba Samuel, Mustapha Adams, Vivian Nathaniel, Asmau M. Maude, Mohammed Evuti Mahmud, Abd'Gafar Tunde Tihamiyu. (2024). *Microplastics and Nanoplastics in Soil. Phytoremediation in Food Safety*. Taylor and Francis Group. CRC Press. Chapter (8), Pg 10.

DOI:<https://doi.org/10.1202/9781032683751>

ISBN: 9781032683751